

Keep this card with you for 3 months after the last RoACTEMRA dose, since side effects could occur for some time after your last dose of RoACTEMRA.

Doctor's Phone:

Doctor's Name:

Patient's Name:

*Please make sure you also have a list of all your other medicines with you at any visit to a healthcare professional.

Most recent:

Start:

Dates of RoACTEMRA Treatment:



RoACTEMRA Patient Alert Card

This patient alert card contains important safety information that you need to be aware of before and during treatment with RoACTEMRA.

- Show this card to any doctor involved in your care
- Read the RoACTEMRA package leaflet for more information



Infections

RoACTEMRA increases the risk of getting infections, which can become serious if not treated.

- You should not be treated with RoACTEMRA if you have a severe infection
- You should have been screened for tuberculosis prior to treatment with RoACTEMRA
- Talk to your doctor about any vaccinations you may need before you start treatment with RoACTEMRA
- If you have an infection of any kind (even a head cold) at the time of your next treatment, the infusion should be delayed until you are feeling better
- Seek medical attention immediately if you develop symptoms such as fever, persistent cough, weight loss, throat pain or soreness, wheezing, red or swollen skin blisters, tears, wounds, severe weakness or tiredness

Complications of Diverticulitis

Patients using RoACTEMRA may develop complications of diverticulitis, which can become serious if not treated.

- Seek medical attention immediately if you develop stomach pain, colic or notice blood in your stools